

# **Disorder Effects on Massive Dirac Fermions**

---

**Qian Niu**

**The University of Texas at Austin**

**ICQM, Peking University**

# Collaborators

---

- Shengyuan Yang, Zhenhua Qiao (The University of Texas at Austin)
- Yugui Yao, Junren Shi (Institute of Physics, Peking University)

# Outline

---

- Introduction
- Scattering Universality Classes: implications on side jump Hall conductivity
- Spin Flip Scattering on Massive Dirac Fermions: Weak Scattering Regime
- Spin Flip Scattering on Massive Dirac Fermions: Strong Scattering Regime



# Massive Dirac Fermion: Introduction

---

# Massive Dirac Fermion: Semiclassical view

---

a wavepacket in the upper band



Has a magnetic moment, and feels a Berry curvature which leads to spin Hall effect and spin Nernst effect


$$\hbar \dot{\mathbf{k}}_c = -e\mathbf{E} - \frac{e}{c} \frac{\hbar \mathbf{k}_c}{\epsilon m} \times \mathbf{B},$$

$$\dot{\mathbf{r}}_c = \frac{\hbar \mathbf{k}_c}{\epsilon m} + \frac{e}{\hbar} \left( \mathbf{E} \times \mathbf{F} + \mathbf{B} \cdot \mathbf{F} \frac{\hbar \mathbf{k}_c}{\epsilon m c} \right),$$

C.-P. Chuu, M.-C. Chang and Q. Niu, 2010

# Massless Dirac Fermion

Graphene:



Topological insulator:



Bi<sub>2</sub>Te<sub>3</sub> surface state


$$\mathcal{H}(\mathbf{k}) = v\mathbf{k} \cdot \boldsymbol{\sigma}$$

# Desire for Opening a Gap


---

1. Gap is essential for any logical device applications.
2. Nonzero Berry curvature leads to interesting physics.

staggered potential on graphene



Magnetically doped TI






Y. L. Chen et al., 2010

# Massive Dirac Fermions from TI thin film

Hai Zhou Lu...SQ Shen:  
PHYSICAL REVIEW B  
**81**, 115407 2010

Yi Zhang ... QiKunXue:  
[Nature Physics 6,](#)  
[584-588 \(2010\)](#)






# Berry Curvature and Hall Transport

Intrinsic contribution to Hall conductivity:

$$\sigma_{xy}^{\text{int}} = -e^2 \int \frac{d\mathbf{k}}{(2\pi)^2} f_{\mathbf{k}} \Omega^z(\mathbf{k}) = -\frac{e^2}{4\pi} \frac{\Delta}{\mu} \quad (\mu > \Delta)$$

Valley Hall effect in graphene:



Di Xiao, Wang Yao, and Qian Niu, 2007

Disorder Scattering, interplayed with Berry curvature, provides another important contribution.

# Outline

- What is a Massive Dirac Fermion?

# Scattering in Spin-Orbit Coupled System

---

- Disorder scattering strongly affects transport properties.
- Scattering has various origins: impurity, dislocation, surface roughness, phonon, magnon, ...

For spin-orbit coupled system, the effects of scattering sensitively depends on its spin structure.

Three universality classes of scattering for 2D systems:

$$\text{Class A} \quad \hat{V} = V^o \hat{1},$$

$$\text{Class B} \quad \hat{V} = V^o \hat{\sigma}_z,$$

$$\text{Class C} \quad \hat{V} = V^o \hat{\sigma}_{\pm} / \sqrt{2},$$

# Scattering Universality Classes

---

Class A	$\hat{V} = V^o \hat{1},$	
Class B	$\hat{V} = V^o \hat{\sigma}_z,$	
Class C	$\hat{V} = V^o \hat{\sigma}_{\pm} / \sqrt{2},$	Spin-flip scattering

For ferromagnet:

Class A: normal impurity, phonon ...

Class B: magnetic impurity in the mean magnetization direction ...

Class C: magnon ...


For graphene:

Class A: on-site impurity, acoustic phonon ...

Class C: optical phonon ...

# Scattering Universality Classes

Different scattering class leads to different anomalous Hall conductivity.



Intrinsic + side jumps

It was believed that side jump Hall conductivity does not depend on the strength and range of scattering.

We found: side jump Hall conductivity depends strongly on the symmetry type of scattering.

For mixed type scatterings, the result depends on the relative strength of different types of scatterings

# Outline

---

- What is a Massive Dirac Fermion?
- Scattering Universality Classes
- **Spin Flip Scattering: Weak Scattering Regime**
- Spin Flip Scattering: Strong Scattering Regime

# Berry Curvature Density Berry curvature per unit energy interval

---

- Generalize the Berry curvature concept to disordered systems

For each eigenstate  $\alpha$ , define Berry curvature for this state as

$$\Omega_{\alpha} \equiv - \sum_{\beta \neq \alpha} \frac{2 \operatorname{Im} \langle \alpha | \nabla | \beta \rangle \langle \beta | \nabla | \alpha \rangle}{(\omega_{\beta} - \omega_{\alpha})^2}$$

Its spectral distribution is described by density of Berry curvature, which is defined as

$$\Omega(\varepsilon) = \sum_{\alpha} \Omega_{\alpha} \delta(\varepsilon - \omega_{\alpha})$$


The density of Berry curvature completely determines the Hall conductivity

$$\sigma_{xy} = -e^2 \int d\varepsilon \langle \Omega(\varepsilon) \rangle_c f(\varepsilon)$$


Sum rule: integral of Berry curvature density between mobility edges is an integer

# Berry Curvature Compression

Perturbative calculation




With spin-flip scattering, Berry curvature distribution is singularly compressed to band edges.



Flat Hall plateaus even for partially filled bands


# Berry Curvature Compression

Numerical evaluation from Kubo formula with disorder average,



These results agree with perturbative results in the thermodynamic limit.

# Density of States and Spin Polarization



DOS and spin polarization are greatly enhanced at band edges, which can be explained by a self energy with a real part diverging at band edges.

$$\text{Re}\Sigma^R(\varepsilon) = -\frac{n_{\text{dis}}V^2}{2\pi v^2} \ln \left| \frac{\varepsilon^2 - \varepsilon_c^2}{\varepsilon^2 - \Delta^2} \right| (\varepsilon\sigma_0 - \Delta\sigma_z)$$

# Outline


---

- What is a Massive Dirac Fermion?
- Scattering Universality Classes
- Spin Flip type: Weak Scattering Regime
- **Spin Flip type: Strong Scattering Regime**

# Lattice Model Hamiltonian

$$H = \lambda(\sigma_x \sin k_x + \sigma_y \sin k_y) + (\cos k_x + \cos k_y + \Delta)\sigma_z$$

In the non-interacting Dirac model, half integer quantum Hall conductance is not possible. So this means there must be other contributions from the Brillouin zone to add or cancel this contribution, so that the total Hall conductance becomes an integer.



(1) Dirac contribution


(2) Conventional contribution

$k_x$


$\Omega$

# Averaged Two-Terminal Conductance

Landau-Buttiker formula + Green's function method




Spin-flip disorder




Spin-independent disorder

# Real Space Calculation of Finite-Size Kubo Formula




Similar character as that of two-terminal conductance


# Evolution of Berry Curvature Density




The Dirac component of the Berry Curvature Density of the electron regime moves towards the hole side, and vice versa (green arrow). While the conventional component moves towards the opposite direction (red arrow).

# Evolution of Berry Curvature Density






# Evolution of Berry Curvature Density (Schematic)



# Phase Diagram (preliminary result)



Anderson Insulator I:  $\text{DOS} \neq 0$ .

Anderson Insulator II:  $\text{DOS} = 0$ .

Metallic I:  $\sigma_{xy} \neq 0$  and  $\sigma_{xx} \neq 0$ .

Metallic II:  $\sigma_{xy} = 0.5e^2/h$  and  $\sigma_{xx}$  is a constant.

Quantum Hall Insulator:  $\sigma_{xy} = e^2/h$  and  $\sigma_{xx} = 0$ .

# Conclusions

---

- What is a Massive Dirac Fermion?
- Scattering Universality Classes
- Spin Flip Scattering on Massive Dirac Fermions: Weak Scattering Regime
- Spin Flip Scattering on Massive Dirac Fermions: Strong Scattering Regime

---

**Thank You!**