

Outlines

- I. *Electromagnetic wave multiple scattering theory* for arrays of spherical particles
- II. The use of particles in construction of *complete band gap photonic crystals*
- III. The use of colloidal crystals in construction of *quasi-3D plasmonic crystals*
- IV. *Light tunneling through* 2D ordered dense array of metal nanoshells
- V. Summary

1. Electromagnetic wave (EM) multiple scattering theory (MST)

- V Development of an analog of KKR method for electron in solid, for photons propagating in arrays of spherical particles;
- V Construction of the software programs;
- V Various implementations of the theory.

W.Y. Zhang et al., PRL, 84, 2853 (2000);

Z.L. Wang et al., PRB 64, 113108 (2001);


Z.L. Wang et al., PRE 67, 016612 (2003);

H. Chen et al., J. Phys.: Condens. Matter 16, 741 (2004);

Typical Approaches for PC in visible/near IR


via microfabrication

S. -Y. Lin, *Nature* 394, (1998) 251;
S. Noda et al., *Science* 289, (2000) 604




via holographic lithography

M. Campbell et al., *Nature* 404 (2000) 53.



via self-assembly

A. Blanco et al., *Nature* 405 (2000) 437.
Y. A. Vlasov et al., *Nature* 414 (2001) 289.




The features of EM MST

- ü Applicable to photonic crystals composed of dielectric *spheres*, either in *simple* lattices, or in *complex* lattice, i.e., diamond lattice;
- ü Much *improvement in precision* as compared to plane-wave-expansion method
- ü Applicable to photonic crystals consisting of core/shell *composite spheres*
- ü Applicable to spheres with dielectric *dispersion or absorption* without CPU time increase.
- ü **Disadvantage:** not applicable to crystals with point or line defects


Specific implementations of EM MST

- ü We clarified the photonic band structure of diamond photonic crystals composed of dielectric spheres;
- ü In contrast to the well known prediction made in *PRL* 65 (1990) 3152 by Ho et al., **the 2nd-3rd** band gap is found to be no larger than 2%
- ü We designed two kinds photonic crystals, both with a complete band gap.

Photonic band structure of diamond photonic crystal of dielectric spheres, calculated using different methods



Based on plane-wave expansion
Ho et al., Phys. Rev. Lett. 65 (1990) 3152



Based on **EM MST**

Design of photonic crystals with complete band gap





The band structure of a photonic crystal is related to crystal symmetry, and are also closely related to the optical properties of the building blocks of the crystals.




A) photonic crystals made of **hollow dielectric** spheres **in a diamond lattice**;

B) photonic crystals made of **metal-coated spheres** in a **face-centered-cubic lattice**



(SEM of silica opal)









- B.T. Holland *et al.*, *Science*, 281 (1998) 538
A.A. Zakhidov *et al.*, *Science*, 282 (1998) 897
S.A. Johnson *et al.*, *Science*, 283 (1998) 963
A. Blanco *et al.*, *Nature*, 405 (2000) 437.
Y.A. Vlasov *et al.*, *Nature*, 414 (2001) 289.
R.F. Service, *Science*, 295 (2002) 2399.
P. Lodahl *et al.*, *Nature* 430 (2004) 654.

Solution A: Photonic crystals made of hollow dielectric spheres in a diamond lattice


$$U \propto \frac{1}{\epsilon}$$






Opening a true gap in diamond photonic crystals using hollow dielectric spheres




Comparison of the gap size in different structures



True gap diamond lattice photonic crystals composed of hollow dielectric spheres




Experiment




Calculations


Assuming that hollow dielectric spheres are arranged in air in a diamond lattice.



Silica beads in diamond lattice




(PS + silica spheres)




(silica spheres)

[111]



(PS + silica spheres)




(silica spheres)


[100]

F.G.S. et al. Adv. Mater. 14(2002), 1144

Solution B: photonic crystals made of metal-coated spheres in a face-centered-cubic lattice





Z. L. Wang, et al., **Phys. Rev. B** 64, 113108 (2001) W. Y. Zhang, et al., **Phys. Rev. Lett.** 84, 2853 (2000).




Growth of onion-structure particles via coating on supporting spheres

Dielectric/Metallic




G. Kaltenpoth et al., *Adv. Mater.*
15 (2003) 1113.

Dielectric/Dielectric






Results from our group



E. Prodan, et al. *Science* **302** (2003) 419



Ordered arrays of silver hollow spheres via colloidal crystal templating




Z. Chen et al., *Adv. Mater.* **16**, 417 (2004)

Questions to be solved:

- ü How to obtain a **non-close-packing of metallic nanoshell** array, with controllable separations?
- ü How to deal with challenges in growth of the metallic nanoshells, such as **rough surface, low coverage, and fragility**?




G. Kaltenpoth et al., *Adv. Mater.* **15** (2003) 1113.






E. Prodan, et al. *Science* **302** (2003) 419

Step one: Non-closely packed colloidal crystals




Schematic




SEM images of a non-close-packing of silica colloidal crystals

Step two and three: Gold nanoshells interconnected with gold nanotubes by double templating




Schematic



SEM images of an inverse polymer opal

Au nanoshells interconnected with Au nanotubes via electroless plating in inverse poly opal






SEM images of a non-close-packing gold nanoshells

W. Dong et al., *Adv. Mater.* **18**, 755 (2006).



The specular optical reflection of a monolayer gold nanoshells



experimental




calculations



**3. The use of colloidal crystals in construction
of *quasi-3D plasmonic crystals***


Preparation of quasi-3D plasmonic crystals



(2D colloidal crystal covered with a thin gold layer)

P. Zhan et al., *Adv. Mater.* **18** (2006) 1612.

Anomalous light transmission



Scalable optical properties




$$\frac{\lambda_{r1}}{n_1} = \frac{\lambda_{r2}}{n_2}$$

$n = 1.45$;

$n = 1.59$)







00 800




Critical dependence on metal film thickness



Assumption: a transition between the excitation of localized SPs and SPP.

Large single domain of the template produced via self-assembly under capillary forces in micro-channels






2D square lattice colloidal crystals




Questions to be solved:
short-range order, high
density of defects

colloidal epitaxy






A. van Blaaderen et al., *Nature*
385, 321 (1997)

self-assembly on 1D patterns




K. Lin et al., *Phys. Rev. Lett.* **85**,
1770 (2000).







Only for α
controlled in the
vicinity of 45°



By controlling the angle α in the vicinity of 45° , a highly ordered 2D square-lattice colloidal crystal with a minimum density of defects can be self-organized.



One of the possible applications

**Quasi-3D plasmonic crystal sensors
with an enhanced sensitivity**

Y. Y. Li et al., to be published.

4. Light tunneling through 2D ordered array of metal nanoshells

C. J. Tang et al., to be submitted.

Basic optical properties and applications of metal films with hole or slit arrays

Å *Ultrafast response*

M. Tong et al., *PRL* 100, 056808 (2008)

Å *Strong modification of nonlinear optical response*

J. A. H. van Nieuwstadt et al., *PRL* 97, 146102 (2006)

Å *Interplane coupling*



X. Fang et al., *PRL* 99, 066805 (2007)

Å *Sensing, data storage, light beam collimation, and extraction from LEDs*

.....

Å *Question about the mechanism associated with enhanced transmittance*

Strong influence of hole shape on extraordinary transmission (experiment)






K.J.K. Koerkamp et al., ***PRL*** 92, 183901 (2004).
(U. Twente, The Netherlands)

Different resonances involved (theory):

- surface plasmon resonances due to the periodicity.
- waveguide resonances in the nanohole with

Terahertz electromagnetic wave transparency




terahertz transparency can be realized through the shape resonance only, when the hole shape is strongly asymmetric.

J.W. Lee et al., PRL **99**, 137401 (2007)

V Sphere surface mode

V Cavity mode



5. Summary

- We developed an efficient MST for EM wave in structures composed of ordered array of spheres
- We designed two types of photonic crystals with true gap using spheres as building blocks.
- We revealed the enhanced optical transmission and other properties of quasi-3D plasmonic crystals.
- We predicted various tunneling channels for light through 2D dense array of metallic

Collaborators

Prof. C. T. Chan, Prof. Ping Sheng

The Hong Kong University of Science & Technology

Prof. J. Zi

Fudan University

Prof. G. Sun

The Institute of Physics, CAS

Prof. X. Y. Jiang

Institute of Microsystem and Information Technology, CAS

Prof. W.Y. Zhang, Prof. S.N. Zhu, Prof. Y.Y. Zhu, Prof. N.B. Ming

Nanjing University

Acknowledgement

Â Distinguished Youth Foundation of NSFC

Â State Key Program for Basic Research of China

Â Key Program of NSFC

Thanks for your attention.